

Agromet Advisory Service Bulletin for Pathanamthitta District (Issued jointly by AMFU, RARS Vellayani, Kerala Agricultural University& India Meteorological Department) Based on Ensemble Weather forecast by IMD



Bulletin No. PTA-AAS-16/2025, (32)

Dated: 22.04.2025

| WEATHER FORECAST FOR THE NEXT FIVE DAYS | | | | | | |
|---|---|------------|------------|------------|------------|--|
| Weather parameters | 23/04/2025 | 24/04/2025 | 25/04/2025 | 26/04/2025 | 27/04/2025 | |
| Rainfall (mm) | 0.3 | 0.4 | 3 | 0.4 | 3 | |
| Maximum temperature(°C) | 35 | 35 | 35 | 35 | 35 | |
| Minimum temperature (°C) | 25 | 25 | 25 | 25 | 25 | |
| Maximum Relative Humidity (%) | 92 | 92 | 92 | 92 | 92 | |
| Minimum Relative Humidity (%) | 60 | 60 | 60 | 60 | 60 | |
| Wind speed (kmph) | 2 | 3 | 3 | 3 | 2 | |
| Wind direction (deg) | 270 | 270 | 270 | 270 | 270 | |
| Total cloud cover (octa) | 8 | 8 | 8 | 8 | 7 | |
| WEATHER SUMMERY ALERT | There is a chance of very light rainfall in isolated places of the district. Cumulatively above normal rainfall is likely over kerala. | | | | | |
| GENERAL ADVISORY | Emergency measures should be taken to protect humans, plants, birds, and animals from extreme heat during the summer. To prevent dehydration, drink plenty of water at regular intervals. Due to increased evaporation and persistent dry conditions, ensure timely irrigation for | | | | | |

dehydration, drink plenty of water at regular intervals. Due to increased evaporation and persistent dry conditions, ensure timely irrigation for crops and maintain adequate soil moisture levels.

| AGRO-METEOROLOGICAL ADVISORIES | | | | | |
|--------------------------------|---|--|--|--|--|
| Сгор | Pest/ Disease | Advisory | | | |
| Brinjal | Fruit borer | To control fruit borer infestation in brinjal, remove and destroy the infested plant parts along with the pests. In case of severe infestation, spray Chlorantraniliprole at a concentration of 3 ml per 10 liters of water. | | | |
| Banana | Leaf-eating caterpillar | Leaf-eating caterpillars may infest banana plants. To control them, destroy the affected leaves along with the eggs, larvae, and pupae. Additionally, spray neem seed extract at a 5% concentration. | | | |
| Cucurbitaceous Vegetables | Downy mildew | There is a chance of downy mildew attack in cucurbitaceous vegetables. Spray 2.5gm Mancozeb per one litre of water under the leaves. | | | |
| Animal Husbandry | During periods of overcast or humid weather, avoid bathing the cow or washing the shed, as this can make the heat feel more intense. To improve comfort in the cattle shed, ensure good air circulation by using fans if necessary, keeping the sides open, and, if possible, increasing the roof height. Also, make sure that clean drinking water is readily available both in the shed and in the grazing areas. | | | | |