



Agromet Advisory Service Bulletin for Thiruvananthapuram District

(Issued jointly by AMFU, RARS Vellayani, Kerala Agricultural University &

India Meteorological Department)

Based on Ensemble Weather forecast by IMD



Bulletin No. TVM-AAS- 33/2026, (66)

Dated: 18.08.2026

SIGNIFICANT WEATHER FOR THE PRECEEDING WEEK

Rainfall(mm)	Max Temperature(°C)	Min Temperature(°C)	Relative Humidity (%)	Evaporation(mm)
6.8	31.0-31.4	23.7-24.0	73-95	3.2-4.8

WEATHER FORECAST FOR THE NEXT FIVE DAYS

Weather parameters	19/08/2026	20/08/2026	21/08/2026	22/08/2026	23/08/2026
Rainfall (mm)	1	2	2	1	0.5
Maximum temperature(°C)	32	32	33	33	32
Minimum temperature (°C)	25	25	25	25	25
Maximum Relative Humidity (%)	85	85	85	85	85
Minimum Relative Humidity (%)	80	80	80	80	80
Wind speed (kmph)	6	5	3	3	3
Wind direction (deg)	290	290	290	290	290
Total cloud cover (octa)	3	4	3	4	4

There is a chance of light rainfall in the district.

WEATHER SUMMERY ALERT

GENERAL ADVISORY

- Harvest mature fruits and vegetables during suitable dry periods and keep harvested produce in clean, dry and well-ventilated places.
- In low-lying fields, monitor water levels regularly and take timely measures to drain excess water from the crop root zone.
- Avoid unnecessary field operations when the soil is excessively wet to prevent soil compaction and damage to the crop root system.

AGRO-METEOROLOGICAL ADVISORIES

Rice	➤ Rice bug infestation is commonly observed from the flowering to maturity stages of the crop. ➤ As a precautionary measure, keep the field and bunds free from weeds and grasses. If rice bug infestation is observed, spray fish-jaggery extract @ 20 ml per litre of water. ➤ In case of severe infestation, spray cypermethrin 10% EC @ 0.5 ml per litre of water. ➤ As rice bug infestation coincides with the flowering stage, insecticide application should preferably be carried out before 9:00 AM or after 3:00 PM to minimize adverse effects on pollination.
Banana	➤ Leaf-eating caterpillars are commonly observed feeding on tender banana leaves. They can be controlled by spraying <i>Beauveria bassiana</i> @ 20 g per litre of water. ➤ In case of severe infestation, spray flubendiamide 39.35 SC @ 2 ml per 10

	litres of water or chlorantraniliprole 18.5 SC @ 3 ml per 10 litres of water.
Vegetables	➤ In vegetable crops, the earliest characteristic symptom of magnesium deficiency is interveinal chlorosis (yellowing between the veins) of older leaves. ➤ Magnesium deficiency can be corrected by spraying magnesium sulphate @ 2–5 g per litre of water. ➤ Do not mix magnesium sulphate with potassium-containing fertilizers, as they may be incompatible.